

Influence of Ischemic Pre- and Post-Conditioning on Cardiac Expression of Calcium-Sensing Receptor

Dyukova E., Schreckenberger R., Sitdikova G., Schlüter K.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2016, Springer Science+Business Media New York. Ischemic heart disease is a common cause of patients' death worldwide. Recently, cardiac pre- and post-conditioning (IPC, IPoC) were identified to reduce infarct size. Nevertheless, not only infarct size but also post-infarct remodelling is critical for the long-term enhancing effect. Calcium-sensing receptors (CaSRs) signalling was shown to be involved in IPC and IPoC in the heart. This study aims to clarify CaSRs expression after ischemia-reperfusion injury (I/R), IPC and IPoC. Experiments were performed on adult Wistar rats with left anterior descending coronary artery (LAD) occlusion. Troponin I (TnI) levels were measured in plasma of all animals to quantify the infarct size. Sham-operated animals, rats with I/R, IPC, and IPoC were compared. Left and right ventricular tissue samples from these groups were collected for qRT-PCR analysis. CaSR expression was enhanced in rats with I/R and IPC. Its increase after IPoC was not pronounced. In contrast, left ventricles (LV) showed decreased CaSR expression in rat hearts after I/R, IPC, and IPoC. Data suggest differences in CaSR regulation between LV and RV. Enhanced CaSR expression in RV was observed in tissue with small infarct size.

<http://dx.doi.org/10.1007/s12668-016-0316-8>

Keywords

Calcium-sensing receptor, Ischemia-reperfusion injury, Post-conditioning, Pre-conditioning

References

- [1] Frank, A., Bonney, M., Bonney, S., Weitzel, L., Koeppen, M., Eckle, T. (2012). Myocardial ischemia reperfusion injury: from basic science to clinical bedside. *Seminars in Cardiothoracic and Vascular Anesthesia*, 16, 123-132. doi:10.1177/1089253211436350.
- [2] Kapur, N. K., Qiao, X., Paruchuri, V., Morine, K. J., Syed, W., Dow, S., et al. (2015). Mechanical pre-conditioning with acute circulatory support before reperfusion limits infarct size in acute myocardial infarction. *JACC Heart Failure*, 3, 873-882. doi:10.1016/j.jchf.2015.06.010.
- [3] Thuny, F., Lairez, O., Roubille, F., Mewton, N., Rioufol, G., Sportouch, C., et al. (2012). Post-conditioning reduces infarct size and edema in patients with ST-segment elevation myocardial infarction. *Journal of the American College of Cardiology*, 59, 2175-2181. doi:10.1016/j.jacc.2012.03.026.
- [4] Gan, R., Hu, G., Zhao, Y., Li, H., Jin, Z., Ren, H., et al. (2012). Post-conditioning protecting rat cardiomyocytes from apoptosis via attenuating calcium-sensing receptor-induced endo(sarco)plasmic reticulum stress. *Molecular and Cellular Biochemistry*, 361, 123-134. doi:10.1007/s11010-011-1096-7.
- [5] Wang, R., Xu, C., Zhao, W., Zhang, J., Cao, K., Yang, B., et al. (2003). Calcium and polyamine regulated calcium-sensing receptor in cardiac tissues. *European Journal of Biochemistry*, 270, 2680-2688.

- [6] Tfelt-Hansen, J., Hansen, J. L., Smajilovic, S., Terwilliger, E. F., Haunso, S., Sheikh, S. P. (2006). Calcium receptor is functionally expressed in rat neonatal ventricular cardiomyocytes. *American Journal of Physiology. Heart and Circulatory Physiology*, 290, 1165–1171. doi:10.1152/ajpheart.00821.2005.
- [7] Smajilovic, S., Hansen, J. L., Christoffersen, T. E., Lewin, E., Sheikh, S. P., Terwilliger, E. F., et al. (2006). Extracellular calcium sensing in rat aortic vascular smooth muscle cells. *Biochemical and Biophysical Research Communications*, 348, 1215–1223. doi:10.1016/j.bbrc.2006.07.192.
- [8] Schreckenber, R., Dyukova, E., Sitdikova, G., Abdallah, Y., Schlüter, K. D. (2015). Mechanisms by which calcium receptor stimulation modifies electromechanical coupling in isolated ventricular cardiomyocytes. *Pflügers Archiv - European Journal of Physiology*, 467, 379–388. doi:10.1007/s00424-014-1498-y.
- [9] Sun, J., & Murphy, E. (2010). Calcium-sensing receptor: a sensor and mediator of ischemic preconditioning in the heart. *American Journal of Physiology Heart and Circulatory Physiology*, 299, H1309–H1317. doi:10.1152/ajpheart.00373.2010.
- [10] Bencsik, P., Kupai, K., Giricz, Z., Görbe, A., Pipis, J., Murlasits, Z., et al. (2010). Role of iNOS and peroxynitrite-matrix metalloproteinase-2 signaling in myocardial late preconditioning in rats. *American Journal of Physiology Heart and Circulatory Physiology*, 299, 512–518.
- [11] Bencsik, P., Pálóczi, J., Kocsis, G. F., Pipis, J., Belec, I., Varga, Z. V., et al. (2014). Moderate inhibition of myocardial matrix metalloproteinases2 by ilomastat is cardioprotective. *Pharmacological Research*, 80, 36–42.